

CytoFish Pre-Treatment Kit

SKU: CPTK-002

Intended Use:

For Research Use Only

Application:

For use of Digestion of Thin Prep Slides in preparation for fluorescent in situ hybridization (FISH) and chromogenic in situ hybridization (CISH).

Composition:

Component	Concentration	Size
HCL	0.01N HCL	250ml
Saline Sodium Citrate Buffer	2X SSC	250ml
Pepsin	Not Applicable	5x 40mg

Storage, Handling, Shelf Life, and Disposal:

Store product excluding Pepsin at 15-30°C, Store Pepsin at -15-20°C; avoid frequent fluctuations in temperature. Expiration date noted on product. Gloves and other protective equipment should be used when handling the kit. Dispose as per local regulations.

Warning and Precaution:

Product does not contain any human or animal components. See Safety Data Sheet (SDS) for more detailed safety and handling information. Do not use expired kit. Avoid cross-contamination. Read instructions for use in full before use.

Limitations:

This product is for research use only. Performance is dependent on sample preparation, sample quality, and proper storage. This product is only for use by trained laboratory professionals.

Contact and Support:

3370 Walden Avenue, Suite 114
Depew, New York, 14043
Phone: (716) 856-3873 or 1-800-715-5880
Fax: (716) 431-3440
info@empiregenomics.com
www.empiregenomics.com

Reagent(s) Not Provided:

- Ethanol
- Wash Buffer 1
- Wash Buffer 2

Equipment Not Provided:

- Coplin Jars
- Water Bath
- Humidity Control Cards in dH₂O
- 22x22mm Glass Coverslip
- Lint-Free Wipe
- Timer
- Pipettes (P10)
- Minifuge
- Vortex

CytoFish Pre-Treatment Protocol:

Prep: Prepare 1 mL of 40mg of Pepsin stock in 1mL H₂O; Prepare 40mL of 2xSSC at room temperature. Prepare 1 container of 100% Ethanol and 1 container of 70% Ethanol. Soak Humidity Control Cards in dH₂O.

Digestion

1. Add 1mL of 40mg/mL Pepsin solution to 40mL of pre-warmed, 37°C 0.01N HCL and mix well.
2. Place slide in 37°C pepsin solution for 8-15 minutes.
 - a. **Time is variable and should be determined by the lab performing the tests.**
3. Wash slides in 2x SSC for 1-2 minutes.
4. Immerse slides in 70% Ethanol for 30 seconds.
5. Immerse slides in 100% Ethanol for 30 seconds.
6. Air Dry slides or place on slide warmer.
7. View digestion under microscope. If adequate digestion is not achieved,

repeat steps 2-6 but in 2-minute pepsin increments.

8. If adequate digestion is achieved, proceed to hybridization protocol.

Hybridization

1. Prepare 10 μ l of probe mix per cellular area (2 μ l of probe + 8 μ l of buffer).
2. Pipette the probe on the processed tissue slide on top of the cellular area and place a 22x22 mm coverslip on that area, taking care that there are no air bubbles.
3. Seal the coverslip with rubber cement; ensure all edges of the coverslip are sealed with the rubber cement.
4. Place pre-soaked Humidity Control Cards in HYBrite/ ThermoBrite. Hybridize the slides on the HYBrite/ ThermoBrite with the following program:

Denaturation: 73°C for 3 minutes

Hybridization 37°C for 16-24 hours

5. After incubation period, carefully remove the rubber cement and coverslip. Proceed to wash steps.
6. Post hybridization wash step 1: Place slide in Wash Buffer 1 (73° C) for 2 minutes, agitating vigorously for the first 10-15 seconds.
7. Post hybridization wash step 2: Place slide in Wash Buffer 2 (room temperature) for 2 minutes, agitating vigorously for the first 10-15 seconds.
8. Wipe the back of the slide dry and allow the rest of the slide to fully dry in the dark. Counterstain with DAPI with Antifade and coverslip with 22x50 mm coverslip. Allow the slide to sit in the dark for 15-20 minutes before viewing under the microscope.